Application No.: 10/811,296

2

Docket No.: 03108/0201072-US0

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Original) N-aryloxypropanolyl-N'-phenethyl urea of general formula 3 wherein R is selected from the group consisting of H, 2, 3 or 4-trifluoromethyl, 2, 3, or 4-chloro, 2, 3, c · 4-bromo, 4-acetyl, 4-propionyl, 4-acetamido, 2,3 or 4 methoxy, 4 nitrile, 2,3 or 4-methyl and 4-formyl and X is S or O.

Formula 3

2. (Original) A compound of formula 3 as claimed in claim 1 selected from the group consisting of.:

3a. N-[2-hydroxy-3-(4-trifluoromethylphenoxy)propyl]-N'-2-phenethy-urea.

3b. N-[2-hydroxy-3-phenoxypropyl]-N'-2-phenethyl-urea

3c. N-[2-hydroxy-3-(3-trifluoromethylphenoxy)propyl]-N'-2-phenethyl-urea

3d. N-[3-(4-chlorophenoxy)-2-hydroxypropyl]-N' 2-phenethyl-urea

3c. N-[3-[4-bromophenoxy)-2-hydroxypropyl]-N'-2-phenethyl-urea

3f. N-[3-(4-acetylphenoxy)- 2-hydroxypropyl]-N'-2-phenethyl-urea

3g. N-[2-hydroxy-3-(4-propionylphenoxy)propyl]-N'-2-phenethyl-urea

3h. N-[3-(4-acetamidophenoxy)- 2-hydroxypropyl]-N'-2-phenethyl-urea

{W:\03108\0201072us0\00366241.DOC *031080201072Us0*}

Application No.: 10/811,296 3 Docket No.: 03108/0201072-US0

3i. N-[2-hydroxy-3(2-methoxyphenoxy)propyl]-N'-2-phenethyl-urea

3j. N-[2-hydroxy-3-(4-methoxyphenoxy)propyl]-N'-2-phenethyl-urea

3k. N-[3-(4-cyanophenoxy)- 2-hydroxypropyl]-N''-2-phenethyl-urea

31. N-[2-hydroxy-3-(2-methylphenoxy)propyl]-N'-2-phenethyl-urea

3m. N-[2-hydroxy-3-(3-methoxyphenoxy)propyl]-N'-2-phenethyl-urea

3n. N-[2-hydroxy-3-(4-trifluoromethylphenoxy)propyl]-N'-2-phenethyl-thiourea

30. N-[2-hydroxy-3-(4-propionylphenoxy)propyl]-N'-2-phenethyl-thiourea

3p. N-[2-hydroxy-3-(4-methoxyphenoxy)propyl]-N'-2-phenethyl-thiourea

3. (Original) A process for the preparation of N-aryloxypropanolyl-N'-pheneth l urea derivatives of the formula 3

Formula 3

wherein X is S or O and R is selected from the group consisting of H, 2, 3 or 4-trifluoromethyl, 2,3, or 4-chloro, 2, 3, or 4-bromo, 4-acetyl, 4-propionyl, 4-acetamido, 2,3 or 4 methoxy, 4 nitrile, 2,3 or 4-methyl and 4- formyl,

the process comprising reacting a substituted phenolic compound with epichlorohydrin in the presence of alkali carbonate to obtain the corresponding phenoxy epoxy propane which is then reacted with ammonium hydroxide to obtain aminoalcohol of formula 1

{W:\03108\0201072us0\00366241.DOC *031080201072US0*}

Application No.: 10/811,296

4

Docket No.: 03108/0201072-US0

Formula 1

wherein R is given above, which is then reacted with a cyanate compound of formula 2 wl erein is oxygen or sulphur

Formula 2

to obtain the compound of formula 3.

- 4. (Previously presented) The process as claimed in claim 3 wherein the alkali carbonate is selected from K_2CO_3 and Na_2CO_3 .
- 5. (Previously presented) The process as claimed in claim 3 wherein the reaction between compound of formula 1 and compound of formula 2 is carried out in an aprotic solvent selected from the group consisting of CH₃CN, CHCl₃, CH₂Cl₂ THF and 1,2-Dichloroethans.
- 6. (Previously presented) The process as claimed in claim 3 wherein the reaction between compound of formula 1 and compound of formula 2 is carried out at a temperature in the range of 15-50°C for a period ranging between 5-18 hrs.
- 7. (Previously presented) The process as claimed in claim 3 wherein the reaction between the compound of formula 1 and compound of formula 2 is carried out in equimolar proportions of compound 1 and compound 2.

{W:\03108\0201072us0\00366241.DOC *031080201072US0*}

Application No.: 10/811,296

5

Docket No.: 03108/02010 2-US0

- 8. (Currently amended) A pharmaceutical composition comprising a compound of formula 3 of claim 1 elaim 2 with one or more conventional additives.
- 9. (Currently amended) A method for the treatment of obesity comprising administering to a subject suffering from obesity, a pharmaceutically effective amount of a compound of formula 3 of claim 1 elaim 2.
- 10. (Previously presented) The method as claimed in claim 9 wherein the compound of formula 3 is administered in the form of a pharmaceutical composition of compound of formula 3 with pharmaceutically acceptable additives.
- 11. (Previously presented) The method as claimed in claim 9 wherein the compound of formula 3 is administered alone or with one or more pharmac outlically acceptable excipients.
- 12. (Currently amended) The method as claimed in claim 16 11 wherein N-[2-bvdroxy-3-[4-trifluoromethylphenoxy]propyl]-N'-2-phenethylurea (3a) shows same activity as Siburramine.
- 13. (Currently amended) The method as claimed in claim 16 H wherein (N -[2-1ydroxy-3-(4-trifluoromethylphenoxy) propyl]-N'-2-phenethyl-urea, showed a decrease of 41.42% n food intake as compared to food intake in the control group.
- 14. (Currently amended) The method as claimed in claim 16 11 wherein (N-[2-1 ydroxy-3-(4- methoxyphenoxy) propyl]-N'-2-phenethyl-urea (compound 3j), showed decrease of 3 .82% in food intake as compared to food intake in the control group.
- 15. (Currently amended) The method as claimed in claim 16 11 wherein (N [2-hydroxy-3-(4- trifluoromethylphenoxy)propyl]-N'-2-phenethyl-thiourea (compound 3n), shawed decrease of 28.4% in food intake as compared to food intake in the control group.
- 16. (New) A method for the treatment of obesity comprising administering to a subject suffering from obesity, a pharmaceutically effective amount of a compound of claim 2.

{W:\03108\0201072us0\00366241.DOC *031080201072Us0*}